

گونه اسکورزونرا اینسیزا از تیره کاسنیان، به عنوان گزارشی جدید از شمال غرب ایران

سیدرضا صفوی

بخش گیاه‌شناسی، موسسه تحقیقات جنگل‌ها و مراتع کشور، سازمان تحقیقات، آموزش و ترویج کشاورزی، ایران؛ (safavi@rifr-ac.ir)

چکیده. *Scorzonera incisa* که گیاهی انحصاری ترکیه محسوب می‌شده است، از استان آذربایجان غربی در شمال غرب ایران جمع‌آوری گردیده و برای اولین بار از کشور گزارش می‌شود؛ این گونه بسیار شبیه به *Scorzonera calyculata* است که در نیمه غربی ایران و شرق عراق پراکنش دارد ولی به دلیل شکل خاص فندقه از آن جدا می‌شود. در این مقاله شرح، تصویر نمونه هرباریومی و نقشه پراکنش این گونه ارائه می‌شود.

واژه‌های کلیدی. آذربایجان غربی، تیره مرکبان، رده‌بندی گیاهی، فلور ایران، لاکتوسه

Scorzonera incisa (Asteraceae), as a new record from NW Iran

Seyed Reza Safavi

Botany Research Division, Research Institute of Forests and Rangelands, Agricultural Research, Education and Extension Organization (AREEO), Tehran, Iran; (safavi@rifr-ac.ir)

Abstract. *Scorzonera incisa*, known as an endemic plant to Turkey, has been collected from West Azarbayegan Province, NW. Iran. The species is reported for the first time from the country; it is very similar to *Scorzonera calyculata*, which is distributed in the western half of Iran and east of Iraq. However, with regard to the particular shape of the achenes of *Scorzonera incisa*, this species could be separated from *Scorzonera incisa*. In this paper, the description, the image of the herbarium sample and the distribution map of this species are presented.

Keywords. Compositae, Iran's flora, Lactuceae, plant taxonomy, West Azarbayegan

INTRODUCTION

The author has worked on the project "Flora of Iran (Asteraceae, Lactuceae)" in the herbarium of the Research Institute of Forests and Rangelands (TARI) since 1998. During a scientific excursion in 2003, a new record species of *Scorzonera* L. was collected from West Azarbayejan Mountains, in NW Iran. Since the genus *Scorzonera* was revised by Rechinger (1977) in the "Flora Iranica", two new species (Safavi, 2006a & 2016) and one new variety to the world (Safavi, 2006b) as well as two species as new records (Dyanat-Nejad *et al.* 1998; Safavi, 2004) have been added to the flora of Iran.

MATERIALS AND METHODS

During the work on the taxonomy of *Lactuceae* tribe in Iran, some specimens of *Scorzonera* were

collected from West Azarbayejan Province in NW Iran and deposited in TARI by the author. After a thorough determination of the specimens, a new record of *Scorzonera* from Iran was recognized. The new record was compared with a large number of specimens belonging to the *Scorzonera* species deposited in the herbaria of TARI and the Iranian Research Institute of Plant Protection (IRAN); it was also compared with the published descriptions of *Scorzonera* species in the flora of Iran and neighboring countries (Chamberlain, 1975; Lipschits, 1964; Rechinger, 1955, 1977; Safavi *et al.* 2013). After a careful consideration, it was concluded that the specimen was a new record for the flora of Iran, belonging to the *Scorzonera incisa* DC.



Fig.1. *Scorzonera incisa* DC. (Safavi & Nikchehreh 86132, TARI).



Fig. 2. *Scorzonera incisa* DC. Achene & Capitule, (Safavi & Nikchehreh 86132, TARI)

RESULTS

New Record to Iran

Scorzonera incisa DC. Iran. West Azarbayejan: Makou, Chaldoran to Khoy, Mahlamlou, 4 Km on the road from Navar to Yekmaleh, N: 38° 54' 09"; E: 44° 14' 09", 2451m., 28 Jun 2003, Rocky slopes, Safavi & Nikchehreh 86132 (TARI) (Figs. 1 & 2).

Erect caulescent perennial 20-40 cm; rootstock cylindrical. Basal leaves dentate to pinnatifid entire, 10-23×2.4-8 cm, stem leaves at most half size of basal leaves. Capitula 1-2-(4) (-many) per stem, (35-) 40-50 mm long. Inner phyllaries 25 mm. Flowers

yellow with dark purple tube. Achenes 12-17 mm, deeply sulcate, with warty wings, sometimes swollen above base (galled) so appearing pyriform and stipitate, glabrous; pappus dirty white, hairs short and plumose or longer, barbellate above and plumose below.

Habitat: Mountainous steppe and Rocky slopes and cliffs, 800 to 2800 m.

Distribution: Subendemic of Turkey, with one record from NW. Iran, An Ir.-Tur. element (Fig 3).




Fig. 3. Distribution of *Scorzonera incisa* DC. in Iran (●)

DISCUSSION AND CONCLUSION

Scorzonera incisa is similar to *Scorzonera calyculata* Boiss., which is distributed in Iran (N., NW., W., S. & C.) and Iraq (Safavi *et al.*, 2013). *S. incisa* belongs to subgen. *Scorzonera* sect. *Incisa* Lipsch. Also, it differs from *S. calyculata* in many features; for instance, achenes 12-17 mm, deeply sulcate, with warty wings, sometimes swollen above base (galled) so appearing pyriform and stipitate, glabrous (not 18 mm, cylindrical-fusiform, glabrous, costate, costa suberose-pallescens, nonstipitate); Pappus dirty white, hairs short and plumose or longer, barbellate above and plumose below (not clear white, almost as long as achene, hairs plumose and scabrous in the upper part).

ACKNOWLEDGMENT

The author wishes to thank Dr. Mohammad Mahmoodi (RIFR, Tehran, Iran) for preparing the map.

REFERENCES

- Chamberlain, D.F.** 1975. *Scorzonera incisa* DC. – In: Davis, P.H. (ed.), Flora of Turkey and East Aegean Islands 5: 639, 644, 645 & 649. – Univ. Press, Edinburgh.
- Dyanat-Nejad, H., Mozaffarian, V. and Safavi, S.R.** 1998. *Scorzonera Koelpinoides* (Asteraceae), a new

record for the flora of Iran. – Iranian J. Bot. 7: 265-267.

Lipschits, S.Yu. 1964. *Scorzonera* L. – In: Bobrov, E.G. and Tzvelev N.N. (eds.), Flora of the USSR 29: 31-144. – Botanical Institute of the Academy of Science of the USSR Press, Moscow, Leningrad.

Rechinger, K.H. 1955. *Scorzonera* L. Symbolae Afghanicae 2: 19-202. – Akademische Druck- u. Verlagsanstalt, Graz.

Rechinger, K.H. 1977. *Scorzonera* L. – In: Rechinger, K.H. (ed.), Flora Iranica 122: 16-79. – Akademische Druck- u. Verlagsanstalt, Graz.

Safavi, S.R. 2004. A new record and an interesting species of the genus *Scorzonera* L. from Iran. – Iranian J. Bot. 10: 159-162.

Safavi, S.R. 2006a. A new species of *Scorzonera* (Asteraceae) from Khorasan, Iran. – Nord. J. Bot. 24: 261-264.

Safavi, S.R. 2006b. Notes on the genera *Scorzonera* L. and *Scolymus* L. (Asteraceae) in Iran. – Iranian J. Bot. 12: 59-62.

Safavi, S.R., Naseh, Y., Jafari, E., Tavakoli, Z. and Heydarnia, N. 2013. *Scorzonera* L. – In: Assadi, M., Maassoumi, A.A., Mozaffarian, V. and Safavi, S.R. (eds.), Flora of Iran (Asteraceae, tribe Cichorieae) 70: 352-442. – Research Institute of Forests & Rangelands Press, Tehran, Iran.

Safavi, S.R. 2016. A new species of *Scorzonera* L. (Asteraceae) from Natanz, Iran. – Iranian J. Bot. 22: 1-5.

How to cite this article:

Safavi, S.R. 2019. *Scorzonera incisa* (Asteraceae), as a new record from NW Iran. – Nova Biol. Reperta 6: 334-337.